

Multigraphs (only) satisfy a weak triangle removal lemma

Asaf Shapira *

Raphael Yuster[†]

Abstract

The triangle removal lemma states that a simple graph with $o(n^3)$ triangles can be made triangle-free by removing $o(n^2)$ edges. It is natural to ask if this widely used result can be extended to multi-graphs (or equivalently, weighted graphs). In this short paper we rule out the possibility of such an extension by showing that there are multi-graphs with only $n^{2+o(1)}$ triangles that are still far from being triangle-free. On the other hand, we show that for some $g(n) = \omega(1)$, if a multi-graph (or weighted graph) has only $g(n)n^2$ triangles then it must be close to being triangle-free. The proof relies on variants of the Ruzsa-Szemerédi theorem [16].

1 Introduction

Motivated by a problem in the theory of extremal hypergraphs, Ruzsa-Szemerédi [16] proved the following two theorems.

Theorem 1 (Ruzsa-Szemerédi [16]) *If G is an n vertex graph from which one should remove at least ϵn^2 edges in order to destroy all triangles, then G contains at least $f(\epsilon)n^3$ triangles.*

Theorem 2 (Ruzsa-Szemerédi [16]) *Suppose $S \subseteq [n]$ is a set of integers containing no 3-term arithmetic progression. Then there is a graph $G = (V, E)$ with $|V| = 6n$ and $|E| = 3n|S|$, whose edges can be (uniquely) partitioned into $n|S|$ edge disjoint triangles. Furthermore, G contains no other triangles.*

These two theorems turned out to be two of the most influential results in extremal combinatorics. First, a simple application of these two theorems gives a short proof of Roth's Theorem [15] stating that a subset of $[n]$ of size ϵn contains a 3-term arithmetic progression. The results in [16] were followed by a long line of investigations leading to the recent hypergraph removal lemmas [8, 12, 14, 20], that also lead to new proofs of Szemerédi's Theorem [18] and some of its extensions.

*School of Mathematics and College of Computing, Georgia Institute of Technology, Atlanta GA, 30332. E-mail: asafico@math.gatech.edu

[†]Department of Mathematics, University of Haifa, Haifa 31905, Israel. E-mail: raphy@math.haifa.ac.il

Besides the above applications to additive number theory and extremal hypergraph theory, which were the original motivation for Theorems 1 and 2, they also turned out to have many additional surprising applications. In particular, these theorems also had applications in extremal combinatorics [7, 4], in the study of probabilistically checkable proofs and analysis of linearity tests [9], in communication complexity [13], as well as in testing monotonicity [6] and testing graph properties [1, 2].

Theorem 1, also known as the *triangle removal lemma*, was originally proved for simple graphs, that is, graphs containing no parallel edges. The proof of Theorem 1 applies the regularity lemma [19], which can only handle graphs with constant edge multiplicity¹. In many applications one thus has to be careful and argue that the graph (or hypergraph) on which one tries to apply Theorem 1 is indeed simple; see [17] for one such example. It is thus natural to ask if the removal lemma also holds for multi-graphs with possibly *unbounded* edge multiplicity. Another way of thinking about this question is whether the removal lemma holds when the edges of a graph have *arbitrary* weights. Note that if we were to identify triangles with their vertex sets, then a simple counter example to such a removal lemma would be to take three vertices, and connect each pair with n^2 edges. In multi-graphs, however, we identify a triangle with its set of edges², so the above example actually has n^6 triangles. We first show that even with this way of counting the number of triangles, the removal lemma does not hold in multi-graphs³.

Theorem 3 *There exists a multi-graph G on n vertices, which contains only $n^2 2^{\sqrt{8 \log n}} = n^{2+o(1)}$ triangles, and yet one should remove n^2 edges from G in order to make it triangle-free.*

We note that the edge multiplicity of the multi-graph we use in the proof of Theorem 3 is $2^{\sqrt{8 \log n}} = n^{o(1)}$, so we see that the removal lemma fails even when the edge multiplicity is sub-linear in the size of the graph.

Observe that if we need to remove n^2 edges from a graph in order to make it triangle-free, then it trivially contains at least n^2 triangles. While Theorem 3 states that this trivial lower bound cannot be substantially improved, we can still ask if a minor improvement is possible. The main motivation is that in some cases (e.g., the original one in [16]) one actually only needs to know that if a graph is far from being triangle free then it contains asymptotically more than n^2 triangles. The following theorem answers this question positively.

Theorem 4 *If G is an n -vertex multi-graph from which one should remove at least n^2 edges in order to destroy all triangles, then G contains $\omega(n^2)$ triangles.*

¹The edge multiplicity of a graph is the maximum number of parallel edges between any pair of vertices.

²Note that in simple graphs there is no difference between identifying a triangle with its edge set or its vertex set.

³It was actually stated, without proof, in some papers (see, e.g., [11]) that the removal lemma holds only in simple graphs, although we are not aware of any proof of this fact.

We note that because the proof of Theorem 4 applies Theorem 1, the improvement we obtain is very minor and gives a lower bound of roughly $n^2(\log^* n)^c$ for some $c > 0$ on the number of triangles in the graph. We also remind the reader that Theorem 4 can also be stated with respect to weighted (simple) graphs rather than multigraphs.

2 The proofs

For completeness we start with the short proof of Theorem 2.

Proof of Theorem 2: We define a 3-partite graph G on vertex sets A , B and C , of sizes n , $2n$ and $3n$ respectively, where we think of the vertices of the sets A , B and C as representing the sets of integers $[n]$, $[2n]$ and $[3n]$. For every $1 \leq i \leq n$ and $s \in S$ we put a triangle $T_{i,s}$ in G containing the vertices $i \in A$, $i + s \in B$ and $i + 2s \in C$. It is easy to see that the above $n|S|$ triangles are edge disjoint, because every edge determines i and s . To see that G does not contain any more triangles, let us observe that G can only contain a triangle with one vertex in each set. If the vertices of this triangle are $a \in A$, $b \in B$ and $c \in C$, then we must have $b = a + s_1$ for some $s_1 \in S$, $c = b + s_2 = a + s_1 + s_2$ for some $s_2 \in S$, and $a = c - 2s_3 = a + s_1 + s_2 - 2s_3$ for some s_3 . This means that $s_1, s_2, s_3 \in S$ form an arithmetic progression, but because S is free of 3-term arithmetic progressions it must be the case that $s_1 = s_2 = s_3$ implying that this triangle is one of the triangles $T_{i,s}$ defined above. ■

For the proof of Theorems 3 we will need to combine Theorem 2 with the following well known result of Behrend [3] that was recently slightly improved by Elkin [5].

Theorem 5 (Behrend [3], Elkin [5]) *For every n , there exists $S \subseteq [n]$ of size $n/2^{\sqrt{8 \log n}} = n^{1-o(1)}$ containing no 3-term arithmetic progression.*

Proof of Theorem 3: Let G' be the graph of Theorem 2 when taking $S \subseteq [n]$ to be a 3AP-free set of size $n/2^{\sqrt{8 \log n}}$ as guaranteed by Theorem 5. Let G be the graph obtained by replacing every edge of G' with $n/|S| = 2^{\sqrt{8 \log n}}$ parallel edges. Observe that as G' contains $n|S|$ edge disjoint triangles, one must remove at least $n|S|$ edges from it in order to make it triangle-free. As G contains $n/|S|$ parallel edges for every edge of G' we infer that one must remove n^2 edges from G in order to make it triangle-free. Finally, as G' contains only $n|S|$ triangles, we infer that G contains only $n|S|(n/|S|)^3 = n^2 2^{2\sqrt{8 \log n}}$ triangles, as needed. ■

Proof of Theorem 4: Given a multi-graph G , let T be the simple graph on the same vertex set that contains an edge (u, v) if and only if G has at most $g^2(n)$ edges connecting u and v for some function $g(n) = \omega(1)$ to be chosen shortly. Let's first consider the case that one needs to remove at

least $\frac{1}{2g^2(n)}n^2$ edges from T in order to make it triangle-free. In this case, by Theorem 1, we know that T contains at least $f(\frac{1}{2g^2(n)})n^3$ triangles. Let us now choose a function $g(n) = \omega(1)$ such that $f(\frac{1}{2g^2(n)})n^3 = \omega(n^2)$. This is clearly possible no matter how fast $f(\epsilon)$ goes to 0 with ϵ . Specifically, given the known bounds on $f(\epsilon)$ in Theorem 1 (see, e.g., [10]), one can take $g(n) = (\log^* n)^c$ for some constant $c > 0$. Fixing this choice of $g(n)$ guarantees that in this case T contains $\omega(n^2)$ triangles and so G contains at least this many triangles as well.

So we can assume that we can remove from T a set of edges E of size $\frac{1}{2g^2(n)}n^2$ and thus make it triangle-free. Let us now remove from G all the edges connecting pairs of vertices that are connected by E in T . Note that we thus remove from G at most $g^2(n) \cdot \frac{1}{2g^2(n)}n^2 \leq n^2/2$ edges, hence the new graph we obtain, let's call it G' , has the property that we should remove at least $n^2/2$ edges from it in order to make it triangle-free. Furthermore, each edge in G' has multiplicity at least $g^2(n)$.

Let T' be the simple graph underlining G' , that is, the graph on the same vertex set, with an edge (u, v) if and only if G' has an edge between u and v . Assume first that T' contains at least $n^2/g(n)$ edges that belong to a triangle. In this case T' contains at least $n^2/3g(n)$ triangles, and as the edge multiplicity of G' is at least $g^2(n)$ this means that G' contains at least $n^2g(n)/3$ triangles. As G' is a subgraph of G we infer that G also contains $n^2g(n)/3$ triangles.

So we can now assume that T' has at most $n^2/g(n)$ edges that belong to a triangle. Let E' be a set of minimal size whose removal from G' makes it triangle-free. Let B denote the set of pairs (u, v) for which E' contains at least one edge connecting u and v , and note that by our assumption on T' we have that $|B| \leq n^2/g(n)$. For each pair of vertices $(u, v) \in B$ let $m_{u,v}$ be the number of edges connecting u and v that belong to E' . We claim that for every (u, v) there are at least $m_{u,v}$ paths of length exactly 2 connecting u and v . Indeed, if G' contains less than $m_{u,v}$ such paths, then we can remove the $m_{u,v}$ edges connecting u and v from E' and replace them by one edge from each of the paths of length 2 connecting u and v . The new set has fewer edges and it still makes G' triangle-free, which contradicts the minimality of E' . We thus conclude that for every pair u, v the graph G' has at least $m_{u,v}^2$ triangles containing u and v . Recall that G' still has the property that one should remove at least $n^2/2$ edges from it in order to make it triangle-free. Therefore we have $\sum m_{u,v} = |E'| \geq n^2/2$. Combining the above facts, and using Cauchy-Schwartz, we conclude that the number of triangles in G' (and so also in G) is at least

$$\sum_{(u,v) \in B} m_{u,v}^2 \geq \frac{1}{|B|} \left(\sum_{(u,v) \in B} m_{u,v} \right)^2 \geq \frac{1}{4|B|} n^4 \geq \frac{1}{4} g(n) n^2 ,$$

thus completing the proof. ■

References

- [1] N. Alon, Testing subgraphs in large graphs, *Random structures and algorithms*, 21 (2002), 359-370.
- [2] N. Alon and A. Shapira, A characterization of easily testable induced subgraphs, *Combinatorics, Probability and Computing*, 15 (2006), 791-805.
- [3] F. A. Behrend, On sets of integers which contain no three terms in arithmetic progression, *Proc. National Academy of Sciences USA* 32 (1946), 331-332.
- [4] L. H. Clark, R. C. Entringer, J. E. McCanna and L. Székely, Extremal problems for local properties of graphs, *Combinatorial Mathematics and Combinatorial Computing* (Palmerston North, 1990). *Australas. J. Combin.* 4 (1991), 25-31.
- [5] M. Elkin, An improved construction of progression-free sets, arxiv.org/abs/0801.4310 .
- [6] E. Fischer, E. Lehman, I. Newman, S. Raskhodnikova, R. Rubinfeld and A. Samorodnitsky, Monotonicity testing over general poset domains, *Proc. of STOC 2002*, 474-483.
- [7] Z. Füredi, The maximum number of edges in a minimal graph of diameter 2, *J. of Graph Theory* 16 (1992), 81-98.
- [8] T. Gowers, Hypergraph regularity and the multidimensional Szemerédi theorem, *Ann. of Math.* Volume 166, Number 3 (2007), 897-946.
- [9] J. Håstad and A. Wigderson, Simple analysis of graph tests for linearity and PCP, *Random Structures and Algorithms*, 22 (2003), 139-160.
- [10] J. Komlós and M. Simonovits, Szemerédi's Regularity Lemma and its applications in graph theory. In: *Combinatorics, Paul Erdős is Eighty*, Vol II (D. Miklós, V. T. Sós, T. Szönyi eds.), János Bolyai Math. Soc., Budapest (1996), 295–352.
- [11] D. Král', O. Serra and L. Vena, A removal lemma for linear systems over finite fields, manuscript 2008.
- [12] B. Nagle, V. Rödl and M. Schacht, The counting lemma for regular k -uniform hypergraphs, *Random Structures and Algorithms* 28 (2006), 113-179.
- [13] P. Pudlák and J. Sgall, An upper bound for a communication game related to time-space tradeoffs, *The mathematics of Paul Erdős, I*, (R. L. Graham and J. Nešetřil, eds.), *Algorithms Combin.*, 13, Springer, Berlin, (1997), 393-399.

- [14] V. Rödl and J. Skokan, Regularity lemma for k -uniform hypergraphs, *Random Structures and Algorithms* 25 (2004), 1-42.
- [15] K.F. Roth, On certain sets of integers, *J. London Math. Soc.* 28 (1953), 104-109.
- [16] I. Ruzsa and E. Szemerédi, Triple systems with no six points carrying three triangles, in *Combinatorics* (Keszthely, 1976), *Coll. Math. Soc. J. Bolyai* 18, Volume II, 939-945.
- [17] A. Shapira, A proof of Green's conjecture regarding the removal properties of sets of linear equations, submitted.
- [18] E. Szemerédi, Integer sets containing no k elements in arithmetic progression, *Acta Arith.* 27 (1975), 299-345.
- [19] E. Szemerédi, Regular partitions of graphs, In: *Proc. Colloque Inter. CNRS* (J. C. Bermond, J. C. Fournier, M. Las Vergnas and D. Sotteau, eds.), 1978, 399-401.
- [20] T. Tao, A variant of the hypergraph removal lemma, *J. Combin. Theory, Ser. A* 113 (2006), 1257-1280.